

**The Government of the Union of Myanmar  
Ministry of Education**

**Department of Higher Education (Lower Myanmar)  
and  
Department of Higher Education (Upper Myanmar)**

**Universities  
Research Journal**

# Universities Research Journal 2008

## Vol. 1, No. 2

### Editorial Board

#### **Editors in Chief**

Prof. Dr. Win Naing, Head of the Department of Geology, University of Yangon

Prof. U Khin Aung Than, Head of the Department of Geology, University of East Yangon

U Ali Akaba Khan (a) U Tin Maung Htwe, Department of Geology, University of Mandalay

Prof. Dr. Zaw Win, Head of the Department of Mathematic, University of Yangon

Dr. Soe Soe Hlaing, Head of Mathematics Department, Yangon Institute of Economics

Prof. Dr. Khin Aye Aye, Head of the Department of Mathematics, University of Mandalay

#### **Editors**

Prof. Dr. Aye Ko Aung, Head of the Department of Geology, University of Dagon

Dr. Chit Sein, Head of the Department of Geology, University of Hinthada

Prof. Ohn Myint, Head of the Department of Geology, University of Maubin

Prof. Dr. Than Than Nu, Head of the Department of Geology, University of Mandalay

Dr. Nyan Win, Head of Geology Department, University of Loikaw

U Hla Myint, Head of Department of Geology, Shwebo Degree College

Assis Lect. Dr Khin Khin Lin, Head of the Department of Geology, University of Bhamo

- Prof. Yi Yi Thein, Head of the Department of Mathematics, University of Mawlamyine**
- Prof. Dr. Win Kyi, Head of the Department of Mathematics, Yangon University of Distance Education**
- Prof. Myint Ohn , Head of the Department of Mathematics, University of Patheingyi**
- Prof. Toe Aung, Head of the Department of Mathematics, University of Taungtha**
- Prof. Dr Yee Myint, Head of Department of Mathematics, University of West Yangon**
- Dr. Hla Hla Kyi, Head of the Department of Mathematics, University of Myeik**
- Prof. Nu Nu Naing, Head of the Department of Mathematics, University of Magway**
- Prof. Nu Nu Sein, Head of the Department of Mathematics, University of Monywa**
- Prof. Dr Hnin Oo, Head of the Department of Mathematics, University of Yadanabon**
- Prof. Dr Kay Thi Tin, Head of the Department of Mathematics, Meiktila Institute of Economics**
- Mai Mae Khin, Head of Department of Mathematics, University of Loikaw**
- Prof. Oo Oo Myint, Head of the Department of Mathematics, University of Lashio**
- Asso. Prof. Dr Than Than Oo, Head of the Department of Mathematics, University of Kyaing Tong**

## Contents

	<b>Page</b>
<b>The Geology of Phongum Razi Ranges Putao District, Kachin State</b> <i>Hla Htay and Aung Khin Soe</i>	1
<b>Preliminary report on the study of the rock units exposed along the Mongla - Kengtung - Tarchileik road section, Eastern Shan State</b> <i>Khin Khin Lin</i>	17
<b>Revision of the Stratigraphy and Age of the Early Devonian Zebingyi Formation, Myanmar</b> <i>Aye Ko Aung</i>	31
<b>New Materials of <i>Stegolophodon</i> (Proboscidea, Mammalia) from the Irrawaddy Formation, Myanmar</b> <i>Chit Sein and Tin Thein</i>	49
<b>New Materials of <i>Tetraconodon malensis</i> (Mammalia, Artiodactyla, Suidae) from the Middle Miocene of Central Myanmar</b> <i>Thaung Htike, Zin Maung Maung Thein and Hnin Hnin Htay</i>	65
<b>Palaeoecological and Palaeogeographical Significance of the Rugose Corals from the Middle Devonian "Maymyo Formation" in Pyinoolwin Township</b> <i>Khaing Khaing San</i>	77
<b>The Study of Folding in Calc-silicate Rocks of Pinle-in Area, Mandalay Division</b> <i>Saw Ngwe Khaing</i>	89
<b>A Study of Clay Minerals in the Maubin Area</b> <i>Ohn Myint, Thura Aung, Swe Zin Thant, Nay Soe, Htet Htet Aung, Zizawar Win Naing, Myo Min and Htin Lynn Aung</i>	103
<b>Tin-tungsten Mineralization of Zingyaik-Kadaik Area, Paung Township, Mon State</b> <i>Than Htoo Aung</i>	119

	<b>Page</b>
An Appraisal for the Mineral Assemblages to define Mineral Isograds on Paragneiss from Latha-Phayargnokto Area, Patheingyi Township, Mandalay Division, Myanmar <i>Zaw Win Ko and Win Min Soe</i>	133
Petrology of Devonian Strata in Naungcho Area, Naungcho Township <i>Thein Htike Swe, Yan Naing Htun and Nyan Win</i>	147
Microbialites in the Thitsipin Limestone, Linwe-Kyauktaw Area, Yengan Township, Southern Shan State <i>Khin Khin Lin</i>	163
Stability of Two-Level Difference Method for PDE <i>Soe Soe Hlaing</i>	173
Finite Difference Method for Elliptic Partial Differential Equation <i>Win Win Ei</i>	193
Numerical Approximations for Burgers Equation <i>Su Su Aung</i>	203
Use of Finite Element Methods for Dirichlet Boundary Value Problem <i>Khin Htay Kyi</i>	213
Numerical Method for A Mathematical Model of A Suspension Bridge <i>Soe Soe Aye</i>	231
Propagation Property for Nonlinear Double-Degenerate Parabolic Equation in Multi-Dimension <i>Khin Than Sint</i>	239
A Study on Steady State Drift-Diffusion Model for Semiconductors <i>Khin Thi</i>	247
Numerical Schemes for the Euler Equations <i>Win Win Nwe</i>	261

	<b>Page</b>
Numerical Approximations of One-Dimensional Stationary Drift-Diffusion Model for Semiconductor <i>Cho Sandar</i>	271
Characterizing the Degree Sequences of Signed Cycles and Signed Stars <i>Hla Hla Myint</i>	285
Applications of Eulerian Path and Tour <i>Sandar Myint, Win Win Mar and Yi Myint</i>	295
Spanning Trees with at most $k$ Leaves in a $K_{1,4}$ -free graph <i>Aung Kyaw</i>	307
Arc-Disjoint Path Pair (APP) Problem <i>Nang Kham Maing</i>	323
Characterization of a Bipartite Graph and Its Hamiltonicity <i>Shwin Seinn</i>	337
A Study on Optimal Control of Rice Production in Taungoo Area <i>Aye Ko, Thin Thin Myat and Mi Mi Than</i>	349
Proximal Method for Equilibrium Problems <i>Hla Hla Kyi</i>	367
The Weighted Locally Convex Spaces of Measurable Functions on $N_p$ Family And Continutous Functions on Nachbin Family <i>Saw Marlar Aung and Kyi Htin Paw</i>	373

# **Characterization of a Bipartite Graph and Its Hamiltonicity**

**Shwin Seinn**

## **Abstract**

In this paper, the characterization of a bipartite graph will be first described by using the definitions of cycle, Hamiltonian cycle, odd cycle, and chromatic number in a graph. Then the conditions for a bipartite graph to be Hamiltonian are investigated. Furthermore a necessary condition will be shown for the Hamiltonicity of a complete bipartite graph.

**Key words:** Bipartite Graph, Hamiltonian cycle, Hamiltonicity

## **Introduction**

The most useful object in discrete mathematics is a structure called a graph. Graphs were first introduced in the eighteenth century by the Swiss Mathematician Leonard Euler. One of the reasons for the recent interest in graph theory is its applicability in many diverse fields including computer science, chemistry, electrical engineering, and economics. In the study of graph theory, it is necessary to determine the existence of a cycle in which all the vertices appear exactly once except for the starting and ending vertex that appears twice. Such a cycle is called a Hamiltonian cycle and we say a graph is Hamiltonian if it contains a Hamiltonian cycle. Although there are several kinds of graphs, we are interested in the existence of bipartite graph and its Hamiltonicity.

In this paper, we first describe the basic notations and definitions of graph theory in the first section. Secondly, the characterization of bipartite graph is discussed in the second section. And then, the conditions for a bipartite graph to be Hamiltonian are investigated in the third section. In the last section, a new necessary condition for a bipartite graph to be Hamiltonian is exhibited.

## **Terminology and Notations**

This section describes some definitions, notations and terminology which will be used throughout this paper. Some special definitions that are only relevant in certain sections will also be given where they are needed. Even worse, the terminology in graph theory is not at all standard, so the

reader may find some differences in definitions from any standard textbooks to other ( for example, see Bondy and Murty (1976) and Douglas (2005) ).

A **graph**  $G$  consists of a nonempty finite set  $V(G)$  of elements called **vertices** and a finite set  $E(G)$ , disjoint from  $V(G)$ , of unordered pairs of (not necessarily distinct ) vertices called **edges**. We call  $V(G)$  the vertex set of  $G$  and  $E(G)$  the edge set of  $G$ . We will often write  $G = (V, E)$  which means that  $V$  and  $E$  are the **vertex set** and the **edge set** of  $G$ , respectively. For convenience, we usually denote the edge  $\{u v\}$  ( where  $u$  and  $v$  are vertices of  $G$  ) by  $uv$ . If  $e = uv$  is an edge of  $G$ , then  $e$  is said to **join**  $u$  and  $v$ , and these vertices are said to be **adjacent**. In this case, we also say that  $e$  is adjacent to  $u$  and  $v$  and that  $u$  and  $v$  are **endpoints** of  $e$ . A **loop** is an edge whose two endpoints are the same. A **simple graph** is a graph with no loops and multiple edges. A graph  $G$  is called **complete** if each vertex in  $G$  is connected with every vertex. A complete graph  $G$  on  $n$  vertices is denoted by  $K_n$ . A **subgraph**  $H$  of a graph  $G$  is a graph  $H$  if  $V(H) \subseteq V(G)$  and  $E(H) \subseteq E(G)$ . A graph is said to be **bipartite** if the vertex set  $V$  of  $G$  can be partitioned into two subsets  $X$  and  $Y$  such that each edge connects a vertex in  $X$  and a vertex in  $Y$  or no edges of  $G$  join vertices in the same subset. A **complete bipartite graph** is a bipartite graph with vertex set partitioned into a subset  $X$  of  $m$  elements and a subset  $Y$  of  $n$  elements such that two vertices are connected by an edge if and only if one is in  $X$  and the other is in  $Y$ . A complete bipartite graph on  $m$  and  $n$  vertices is denoted by  $K_{m,n}$ . The **degree** of a vertex  $v$  in a graph  $G$  is the number of edges incident to it and is denoted by  $d(v)$ . A **path** in a graph  $G$  is a sequence of edges that begins at a vertex of a graph and travels along edges of the graph, always connecting pairs of adjacent vertices. The **length** of a path is the number of edges contained in it. If  $u$  and  $v$  are distinct vertices of a graph  $G$  and  $P$  is a path from  $u$  to  $v$ , we say that  $P$  is a  $(u, v)$ -**path**. A graph is said to be **connected** if there is a path between every pair of its distinct vertices. A **cycle** in a graph  $G$  is a path that begins and ends at the same vertex ( i.e., a closed path of non-zero length that does not contain a repeated edge). A **Hamiltonian cycle** is a cycle in a graph  $G$  that contains each vertex of  $G$  exactly once except for the starting and ending vertex that appears twice. A graph is said to be **Hamiltonian** if it contains a Hamiltonian cycle, and otherwise, **non-Hamiltonian**.

## Characterization of a Bipartite Graph

In this section we begin with the definitions of *odd cycle* and *chromatic number* in order to describe the characterization of bipartite graph.

### Definition 1

An odd cycle is a cycle that has an odd number of edges.

### Definition 2

The length of a shortest  $(u, v)$  – path is called the distance from  $u$  to  $v$  and it is denoted by  $d(u, v)$ .

### Definition 3

A  $t$  – coloring of  $G$  is a function  $C : V \rightarrow \{1, 2, \dots, t\}$  such that  $C(u) \neq C(v)$  whenever  $uv \in E$ .

### Definition 4

A graph that has a  $t$  – coloring is said to be  $t$  – colorable.

### Definition 5

The minimum value of  $t$  for which there exists a  $t$  – coloring of  $G$  is called the chromatic number of  $G$  (i.e., the smallest  $t$  such that  $G$  is  $t$  – colorable ), and it is denoted by  $\chi(G)$ .

### Theorem 1

A graph  $G = (V, E)$  is bipartite if and only if it contains no odd cycles.

### Proof:

Suppose that  $G = (V, E)$  is bipartite with bipartition  $(X, Y)$ .

Let  $C = \{v_0 v_1 v_2 \dots v_k v_0\}$  be any cycle of  $G = (V, E)$ . We will show that  $C$  is even cycle. Without loss of generality we may assume that  $v_0 \in X$ . Since  $G = (V, E)$  is bipartite and  $v_0 v_1 \in E$ ,  $v_1 \in Y$ . Similarly  $v_2 \in X$  and in general  $v_{2i} \in X$  and  $v_{2i+1} \in Y$ . Since  $v_0 \in X$ ,  $v_k \in Y$ . Then,  $k = 2i + 1$ , for some  $i$  and it follows that  $C$  is even cycle. Thus, if  $G = (V, E)$  is bipartite, then  $G = (V, E)$  contains no odd cycles.

Conversely, suppose that a connected graph  $G = (V, E)$  contains no odd cycles. Let  $u$  be any vertex of  $G$  and define a partition  $(X, Y)$  of  $V$  by

$$X = \{x \in V : d(u, x) \text{ is even}\},$$

$$Y = \{y \in V : d(u, y) \text{ is odd}\}.$$

We shall show that  $(X, Y)$  is a bipartition of  $G = (V, E)$ .

Suppose that  $v$  and  $w$  are two vertices of  $X$ . Let  $P$  be a shortest  $(u, v)$  – path and  $Q$  be a shortest  $(u, w)$  – path. Denote by  $u_1$  the last vertex common to  $P$  and  $Q$ . Since  $P$  and  $Q$  are shortest paths, the  $(u, u_1)$  – sections of both  $P$  and  $Q$  are shortest  $(u, u_1)$  – paths and, therefore, have the same length. Since the lengths of both  $P$  and  $Q$  are even, the lengths of the  $(u_1, v)$  – section  $P_1$  of  $P$  and the  $(u_1, w)$  – section  $Q_1$  of  $Q$  must have the same parity. It follows that the  $(v, w)$  – path  $P_1^{-1}Q_1$  is of even length. If  $v$  were joined to  $w$ ,  $P_1^{-1}Q_1wv$  would be a cycle of odd length. It contradicts to our assumption. Thus, no two vertices in  $X$  are adjacent; no two vertices in  $Y$  are adjacent. Hence  $G = (V, E)$  is bipartite. ■

## Theorem 2

A graph  $G = (V, E)$  is bipartite if and only if its chromatic number is at most 2.

### Proof:

Suppose that  $G = (V, E)$  is bipartite with bipartition  $(X, Y)$ . Then, a coloring of  $G = (V, E)$  can be obtained by assigning red to each vertex in  $X$  and blue to each vertex in  $Y$ . Since there is no edge from any vertex in  $X$  to any other vertex in  $X$ , and no edge from any vertex in  $Y$  to any other vertex in  $Y$ , no two adjacent vertices can receive the same color. It follows that the chromatic number of the bipartite graph is at most 2.

Conversely, if we have a 2 – coloring  $C$  of a graph  $G = (V, E)$ , let  $X$  be the set of vertices that receive color red and let  $Y$  be the set of vertices that receive color blue. Since adjacent vertices must receive different colors, there is no edge between any two vertices in the same partite set. It follows that  $G = (V, E)$  is bipartite. ■

## Investigations on Hamiltonicity of a Bipartite Graph

In this section we will investigate the existence of Hamiltonian cycle in simple bipartite graph. Now we have question, "which conditions make a bipartite graph to be Hamiltonian?" The answer must be its number of vertices.

### Definition 6

The component of a graph is its maximal connected subgraph.

### Proposition 1

If a graph  $G$  has a set of  $|S| = k$  vertices whose removal from  $G$  results in a graph  $G - S$  which has more than  $k$  connected components, then  $G$  does not have a Hamiltonian cycle.

### Proof:

Suppose that  $G$  has a Hamiltonian cycle. When leaving vertices of a connected component of  $G - S$ , a Hamiltonian cycle can go only to  $S$ . See Figure 1.




Figure 1

Each arrival in  $S$  must be at a different vertex of  $S$ . Hence,  $S$  must have at least as many vertices as  $G - S$  has connected components. It contradicts to the hypothesis of the Proposition. Hence,  $G$  does not have a Hamiltonian cycle. ■

### Theorem 3

If a bipartite graph  $G$  with partite sets  $X$  and  $Y$  is such that  $|X| \neq |Y|$ , then  $G$  is non-Hamiltonian.

**Proof:**

Suppose without loss of generality that  $|X| > |Y|$ . Then, removing the vertices of  $Y$  from  $G$  results in  $|X|$  connected components. So, by Proposition 1,  $G$  does not have a Hamiltonian cycle. Hence,  $G$  is non-Hamiltonian. ■

Consequently, we get the following corollary.

**Corollary 1**

Every bipartite graph having odd number of vertices is non-Hamiltonian. ■

**Example 1**

The graph shown in Figure 2 is the Herschel graph. We will investigate whether it is Hamiltonian.




Figure 2: Herschel Graph

We can review the Herschel graph as a bipartite graph with partite sets,

$$X = \{d, a, y, z, r, q\} \text{ and}$$

$$Y = \{x, s, c, b, p\}, \text{ as shown in Figure 3.}$$




Figure 3

Since  $|X| = 6$  and  $|Y| = 5$ ,  $|X| \neq |Y|$ .

By Theorem 3, the Herschel graph is non-Hamiltonian.

Consequently, one should consider whether the bipartite graph is Hamiltonian if  $|X| = |Y|$ .

The following counterexample shows that although a bipartite graph with partite sets  $X$  and  $Y$  has  $|X| = |Y|$ , it is non-Hamiltonian.

Consider the graph shown in Figure 4.




Figure 4

If the vertices  $v_4$  and  $v_9$  in above graph are removed, then the graph  $G - \{v_4, v_9\}$  has three connected components. Thus, by Proposition 1, the graph does not have Hamiltonian cycle.

The graph of Figure 4 can be transformed into a bipartite graph with  $|X| = |Y|$ , where  $X = \{v_1, v_3, v_5, v_7, v_9, v_{11}, v_{14}, v_{16}, v_{18}, v_{20}\}$  and  $Y = \{v_2, v_4, v_6, v_8, v_{10}, v_{12}, v_{13}, v_{15}, v_{17}, v_{19}\}$ , as shown in Figure 5.




Figure 5

Since  $|X| = 10$  and  $|Y| = 10$ , the redrawing bipartite graph shown above has two equal partite sets.

It is non-Hamiltonian since there does not exist a Hamiltonian cycle. ■

### Necessary Condition for a Complete Bipartite Graph to be Hamiltonian

Our investigations from the previous section are facing with the conflict of which conditions make a bipartite graph to be Hamiltonian. So, the purpose of this section is to show a result we produced which is an agreeable condition for a bipartite graph to be Hamiltonian.

#### Theorem 4

A complete bipartite graph  $K_{m,n}$  is Hamiltonian if and only if  $m = n$ , for all  $m, n \geq 2$ .

#### Proof:

Suppose that a complete bipartite graph  $K_{m,n}$  is Hamiltonian. Then, it must have a Hamiltonian cycle which visits the two partite sets alternately. Therefore, there can be no such cycle unless the two partite sets have the

same number of vertices. If  $m = n = 1$ , it is clear that  $K_{m,n}$  contains no Hamiltonian cycle.

Thus, we get  $m = n$ , for all  $m, n \geq 2$ .

Conversely, suppose that a bipartite graph with partite sets,

$$X = \{x_1, x_2, \dots, x_m\} \text{ and}$$

$$Y = \{y_1, y_2, \dots, y_n\},$$

is complete and its two partite sets have the same number of vertices (i.e.,  $m = n$ ). Then, it is obvious that  $x_1 y_1 x_2 y_2 \dots x_m y_n x_1$  is a Hamiltonian cycle. Hence  $K_{m,n}$  is Hamiltonian. ■

The following example demonstrates Theorem 4.

### Example 2

Consider the bipartite graphs shown in Figures 6 and 7.




Figure 6




Figure 7

The graph of Figure 6 is a complete bipartite graph  $K_{3,3}$  with  $m = n = 3$ .

It satisfies the conditions of Theorem 4. One of its Hamiltonian cycles is

$$x \ l \ v \ w \ u \ o \ x.$$

The graph of Figure 7 is a complete bipartite graph  $K_{4,4}$  with  $m = n = 4$ .

It satisfies the conditions of Theorem 4. One of its Hamiltonian cycles is

$$m \ j \ i \ z \ s \ t \ n \ k \ m.$$

Again, in the following Figure 8, although it is a complete bipartite graph,  $K_{2,3}$ , it is non-Hamiltonian since the numbers of vertices in each partite set are different.




Figure 8

■

### Conclusion

The problem of finding necessary and sufficient conditions for a graph to be Hamiltonian is both interesting and difficult. In this paper, we presented our investigations on the existence of Hamiltonian cycles in bipartite graph. We had exhibited a new necessary condition for a bipartite graph to be Hamiltonian. We firmly believe that there remains a lot to determine in this line of research. Furthermore, one may analyze the Hamiltonicity of bipartite digraphs and the number of Hamiltonian cycles in bipartite digraphs.

### Acknowledgement

I would like to express my sincere gratitude to Pro-Rector Dr Ko Ko Myint of Kyainge Tong University for his kind consent to the submission of this research paper. I'm also deeply indebted to my supervisor, Professor Dr Yi Myint, Head of Department of Mathematics, West Yangon University, for giving me her kindly guidance and suggestions, to the field of Hamiltonicity of Graphs. I also would like to express my unlimited gratitude to Head and Associate Professor Dr Than Than Oo of our department for her helpful comments on doing this research. Finally, I thank to all of my colleagues for their encouragement while doing this research.

## References

- Bondy, J.A and Murty, U.S.R, (1976). *Graph Theory with Applications*, American Elsevier, New York.
- Diestel, R., (2005). *Graph Theory*, Electronic Edition, Springer-Verlag, New York.
- Douglas, B.W., (1999). *Introduction to Graph Theory*, Prentice-Hall of India Private Limited, New Delhi.
- Grossman, J. W., (1990). *Discrete Mathematics*, Macmillan Publi., New York.